FAIRVIEW Fairview-University Medical Center

Hemophilla & Thrombosis Center A collaboration with Children's Hospitals & Clinics -Minneapolis & St. Paul 420 Delaware Street SE, Box 713 Minneapolis, MN 55455 Tel 612-626-6455 Fax 612-625-4955 1-800-688-5252 ext. 66455

January 23, 2002

Commissioner of Social Security PO Box 17703 Baltimore, Maryland 21235-7703

Dear Commissioner:

We submit that the new proposed Listing of Impairments for hemophilia and von Willebrand disorder needs to be reevaluated and modified. Specifically, the issues which are of concern are as follows:

7.03B. and **107.03B.** Hemophilia with spontaneous bleeding despite prophylactic factor replacement, occurring at least 3 times in a consecutive 12 month period.

Comment #1: Prophylaxis is not the universal standard of care. Many adults with severe hemophilia are not on prophylaxis. Prophylaxis is considered to be optimal therapy for children with severe hemophilia, to prevent joint damage and long term disability. However, data collected from the Centers for Disease Control and Prevention indicate that 44% of individuals with severe hemophilia under the age of 18, and 85.5% of patients over the age of 18 are not on prophylaxis. An even greater number of people with moderate hemophilia (>1% factor VIII or FIX) are not on prophylaxis. Therefore, a very significant number of people are on "demand therapy" (treatment of bleeding episodes only at the time of the bleed) and not prophylaxis. Based on the data, we recommend removing the prophylaxis clause from the Listings.

Comment #2: Regarding the joint deformity requirement section of the listing for children, individuals with hemophilia have a different type of joint disease than those with juvenile rheumatoid artiritis. Hemophilia related joint disease is the result of repeated bleeds. Therefore, they need to be listed separately. Alternatively, we urge you to consider whethet a listing in this section is necessary, given that joint bleeds and subsequent arthritic disease are considered to be a complication of hemophilia.

Proposed 113.21E. How Do We Evaluate Episodic Hematological Disorders?

"...h every listing in which we require more than one event (bleeding episode), there must be at least one month between the events (bleeding episodes), in order to ensure that we are evaluating separate episodes."

Comment: People with bleeding disorders may have many separate spontaneous bleeds at various sites occurring during a one month period. Alternatively, an individual could have several separate bleeding episodes in the same muscle or joint during the same month. Therefore, we think this rule needs to be changed.

7.03 and 107.03C. Other hypocoagulable states (such as von Willebrand disorder or thrombocytasthenia) with spontaneous bleeding requiring hospitalization (for 24 hours or mow), occurring at least 3 times in a consecutive 12 month period.

Comment: Individuals with von Willebrand disorder do not regularly require hospitalization to control bleeding episodes, We recommend they not be separated from hemophilia in the new Listings. We recommend the removal of the hospitalization requirement clause.

Sincertyyours

Nigel Key, MD Co-Director Margaret Heisel, MD

largaret Hein

Co-Director

Comprehensive Hemophilia and Thrombosis Center

Fairview-University Medical Center